

Mehrabadi's Conjecture

I have done several joint research with my friend, M.L. Mehrabadi(1969-1998). Unfortunately he passed away in a bus accident. During our joint research about chromatic sum he made an interesting conjecture about strength of a graph [1] and it is open by now. My friend, R. Tusserkani and me decided to call this conjecture, Mehrabadi's conjecture to commemorate him. We will pay \$200 USD as prize to everybody prove or disprove this conjecture.

Let G be a graph, we denote by $\sum_k(G)$ the smallest possible sum among all proper k - colorings of G using natural numbers. The *vertex-chromatic sum* of G , denoted by $\Sigma(G)$, is defined as $\min_{k \geq \chi(G)} \sum_k(G)$. The *vertex-strength* of G denoted by $s(G)$, or briefly by s , is the smallest number s such that $\sum_s(G) = \Sigma(G)$. Clearly, $s(G) \geq \chi(G)$ and equality does not always hold. In fact, for every positive integer k , almost all trees satisfy $s \geq k$ (for more about this concept see [1] and [2]).

Mehrabadi's Conjecture. *For every graph G , we have:*

$$s(G) \leq \lceil \frac{\chi(G) + \Delta(G)}{2} \rceil.$$

After Mehrabadi made his conjecture we have proved the following upper bound for strength of graphs.

Theorem.[1] *For every graph G , we have:*

$$s(G) \leq \lceil \frac{col(G) + \Delta(G)}{2} \rceil,$$

where $col(G)$ is the coloring number of graph G .

References

- [1] H. Hajiabolhassan, M.L. Mehrabadi, and R. Tusserkani, Minimal colorings and strength of graphs, *Discrete Mathematics*, **215**(2000), 265–270.
- [2] H. Hajiabolhassan, M.L. Mehrabadi, and R. Tusserkani, Tabular graphs and chromatic sum, , manuscript(1998).